arXiv:1607.03156v1 [math.AG] 11 Jul 2016

# On the proper push-forward of the characteristic cycle of a constructible sheaf

Takeshi Saito

July 13, 2016

**Abstract**

We study the compatibility with proper push-forward of the characteristic cycles of a constructible complex on a smooth variety over a perfect field.

The characteristic cycle of a constructible complex on a smooth scheme over a perfect field is defined as a cycle on the cotangent bundle [4] supported on the singular support [2]. It is characterized by the Milnor formula [4, (5.15)] for the vanishing cycles defined for morphisms to curves.

We study the compatibility with proper push-forward. First, we formulate Conjecture 1 on the compatibility with proper direct image. We prove it in some cases, for example, morphisms from surfaces to curves under a mild assumption in Theorem 6. We briefly sketch the idea of proofs, which use the global index formula (11) computing the Euler-Poincaré characteristic. For the compatibility Theorem 6, it amounts to prove a conductor formula (14) at each point of the curve. By choosing a point and killing ramification at the other points using Epp's theorem [3], we deduce the conductor formula (14) from the index formula (11). We give a characterization of characteristic cycle in terms of functorialities at the end of the article.

The research was partially supported by JSPS Grants-in-Aid for Scientific Research (A) 26247002.

For the definitions and basic properties of the singular support of a constructible complex on a smooth scheme over a perfect field, we refer to [2] and [4].

Let $k$ be a perfect field and let $\Lambda$ be a finite field of characteristic $\ell$ invertible in $k$. We say that a complex $\mathcal{F}$ of $\Lambda$-modules on the étale site of $X$ is constructible if the cohomology sheaf $\mathcal{H}^q\mathcal{F}$ is constructible for every $q$ and vanishes except for finitely many $q$. More generally, if $\Lambda$ is a finite local ring of residue field $\Lambda_0$ of characteristic $\ell$ invertible in $k$, the singular support and the characteristic cycle of a complex $\mathcal{F}$ of $\Lambda$-modules of finite tor-dimension equals to those of $\mathcal{F} \otimes^L_\Lambda \Lambda_0$.

Let $\mathcal{F}$ be a constructible complex on a smooth scheme $X$ over $k$. The singular support $SS\mathcal{F}$ is defined in [2] as a closed conical subset of the cotangent bundle $T^*X$ By [2, Theorem 1.3 (ii)], every irreducible component $C_a$ of the singular support $SS\mathcal{F} = C = \bigcup_a C_a$ is of dimension $n$. The characteristic cycle $CC\mathcal{F} = \sum_a m_a C_a$ is defined as a linear combination with $\mathbf{Z}$-coefficients in [4, Definition 5.10]. It is characterized by the Milnor formula

$$-\dim \mathrm{tot}\phi_u(\mathcal{F}, f) = (CC\mathcal{F}, df)_{T^*U,u} \tag{1}$$

for morphisms $f\colon U \to Y$ to smooth curves $Y$ defined on an étale neighborhood $U$ of an isolated characteristic point $u$. For more detail on the notation, we refer to [4, Section 5.2].

We say that a constructible complex $\mathcal{F}$ is *locally constant* if every cohomology sheaf $\mathcal{H}^q\mathcal{F}$ is locally constant. In this case, we have

$$CC\mathcal{F} = (-1)^n \mathrm{rank}\ \mathcal{F} \cdot [T^*_X X] \tag{2}$$

where $T^*_X X$ denotes the 0-section and $n = \dim X$ by [4, Lemma 5.11.1]. Assume $\dim X = 1$ and let $U \subset X$ be a dense open subset where $\mathcal{F}$ is locally constant. For a closed point $x \in X$, the Artin conductor $a_x\mathcal{F}$ is defined by

$$a_x\mathcal{F} = \mathrm{rank}\ \mathcal{F}|_U - \mathrm{rank}\ \mathcal{F}_x + \mathrm{Sw}_x\mathcal{F} \tag{3}$$

where $\mathrm{Sw}_x\mathcal{F}$ denotes the alternating sum of the Swan conductor at $x$. Then, by [4, Lemma 5.11.3], we have

$$CC\mathcal{F} = -\Big(\mathrm{rank}\ \mathcal{F} \cdot [T^*_X X] + \sum_{x \in X - U} a_x\mathcal{F} \cdot [T^*_x X]\Big) \tag{4}$$

where $T^*_x X$ denotes the fiber.

To state the compatibility with push-forward, we fix some terminology and notations. We say that a morphism $f\colon X \to Y$ of noetherian scheme is proper (resp. finite) on a closed subset $Z \subset X$ if its restriction $Z \to Y$ is proper (resp. finite) with respect to a closed subscheme structure of $Z \subset X$.

Let $h\colon W \to X$ and $f\colon W \to Y$ be morphisms of smooth schemes over $k$. Let $C \subset X$ be a closed subset such that $f$ is proper on $h^{-1}(C)$ and let $C' = f(h^{-1}(C)) \subset Y$ be the image of $C$ by the algebraic correspondence $X \leftarrow W \to Y$. If $\dim W = \dim X - c$, the intersection theory defines the pull-back and push-forward morphisms

$$CH_\bullet(C) \xrightarrow{h^!} CH_{\bullet - c}(h^{-1}(C)) \xrightarrow{f_*} CH_{\bullet - c}(C'). \tag{5}$$

We call the composition the morphism defined by the algebraic correspondence $X \leftarrow W \to Y$. If every irreducible component of $C$ is of dimension $n$ and if every irreducible component of $C'$ is of dimension $m = n - c$, the morphism (5) defines a morphism $Z_n(C) \to Z_m(C')$ of free abelian groups of cycles.

Let $f\colon X \to Y$ be a morphism of smooth schemes over a field $k$. Assume that every irreducible component of $X$ is of dimension $n$ and that every irreducible component of $Y$ is of dimension $m$. Let $C \subset T^*X$ be a closed conical subset. The intersection $B = C \cap T^*_X X$ with the 0-section regarded as a closed subset of $X$ is called the base of $C$.

Assume that $f\colon X \to Y$ is proper on the base $B$. Then, the morphism $X \times_Y T^*Y \to T^*Y$ induced by $f$ is proper on the inverse image $df^{-1}(C)$ by the canonical morphism $df\colon X \times_Y T^*Y \to T^*X$. Let $f_\circ C \subset T^*Y$ denote the image of $df^{-1}(C)$ by $X \times_Y T^*Y \to T^*Y$. Then $f_\circ C \subset T^*Y$ is a closed conical subset.

By applying the construction of (5) to the algebraic correspondence $T^*X \leftarrow X \times_Y T^*Y \to T^*Y$ and $C \subset T^*X$, we obtain a morphism

$$f_*\colon CH_n(C) \to CH_m(f_\circ C). \tag{6}$$

Further if every irreducible component of $C \subset T^*X$ is of dimension $n$ and if every irreducible component of $f_\circ C \subset T^*Y$ is of dimension $m$, we obtain a morphism

$$f_* \colon Z_n(C) \to Z_m(f_\circ C). \tag{7}$$

Now, let $f \colon X \to Y$ be a morphism of smooth schemes over $k$ and assume that every irreducible component of $Y$ is of dimension $m$. The base $B \subset X$ of the singular support $C = SS\mathcal{F} \subset T^*X$ equals the support of $X$ by [2, Lemma 2.1(i)]. Then, the direct image

$$f_* CC\mathcal{F} \in CH_m(f_\circ C) \tag{8}$$

of the characteristic cycle $CC\mathcal{F}$ is defined by the algebraic correspondence $T^*Y \leftarrow X \times_Y T^*Y \to T^*X$. Further if every irreducible component of $f_\circ C \subset T^*Y$ is of dimension $m$, the direct image

$$f_* CC\mathcal{F} \in Z_m(f_\circ C) \tag{9}$$

is defined as a linear combination of cycles.

**Conjecture 1.** *Let $f \colon X \to Y$ be a morphism of smooth schemes over a perfect field $k$. Assume that every irreducible component of $X$ is of dimension $n$ and that every irreducible component of $Y$ is of dimension $m$. Let $\mathcal{F}$ be a constructible complex on $X$ and $C = SS\mathcal{F}$ be the singular support. Assume that $f$ is proper on the support of $\mathcal{F}$.*

1. *We have*

$$CCRf_*\mathcal{F} = f_* CC\mathcal{F} \tag{10}$$

*in $CH_m(f_\circ C)$.*

2. *In particular, if every irreducible component of $f_\circ C \subset T^*Y$ is of dimension $m$, we have an equality* (10) *of cycles.*

If $Y = \operatorname{Spec} k$ is a point and $X$ is proper over $k$, the equality (10) is nothing but the index formula

$$\chi(X_{\bar{k}}, \mathcal{F}) = (CC\mathcal{F}, T^*_X X)_{T^*X}. \tag{11}$$

This is proved in [4, Theorem 7.13] under the assumption that $X$ is projective. For a closed immersion $i \colon X \to P$ of smooth schemes over $k$, Conjecture 1 holds [4, Lemma 5.13.2]. Hence, for a proper morphism $g \colon P \to Y$ of smooth schemes over $k$, Conjecture 1 for $\mathcal{F}$ and $f = g \circ i \colon X \to Y$ is equivalent to that for $i_*\mathcal{F}$ and $g \colon P \to Y$. For the singular support, an inclusion $SSRf_*\mathcal{F} \subset f_\circ SS\mathcal{F}$ is proved in [2, Theorem 1.4 (ii)].

**Lemma 2.** *Assume that $f$ is finite on the support of $\mathcal{F}$. Then Conjecture* 1.2 *holds.*

*Proof.* We may assume that $k$ is algebraically closed. Since the characteristic cycle is characterized by the Milnor formula, it suffices to show that $f_* CC\mathcal{F}$ satisfies the Milnor formula (1) for $Rf_*\mathcal{F}$.

Let $Z \subset X$ denote the support of $\mathcal{F}$. Let $V \to Y$ be an étale morphism and $g \colon V \to T$ be a morphism to a smooth curve $T$ with isolated characteristic point $v \in V$ with respect to $f_\circ C$. By replacing $Y$ by $V$, we may assume $V = Y$.

By [4, Lemma 3.9.3 (1)⇒(2)] and by the assumption that $Z$ is finite over $Y$, the composition $g \circ f \colon X \to T$ has isolated characteristic points at the inverse image $Z \times_Y v$. Hence, the composition $g \circ f \colon X \to T$ is locally acyclic relatively to $\mathcal{F}$ on a neighborhood of the fiber $X \times_Y v$ except at $Z \times_Y v$ and we have a canonical isomorphism

$$\phi_v(Rf_*\mathcal{F}, g) \to \bigoplus_{u \in Z \times_Y v} \phi_u(\mathcal{F}, g \circ f).$$

Thus by the Milnor formula (1), we have

$$\begin{aligned} -\dim \mathrm{tot}\phi_v(Rf_*\mathcal{F}, g) &= \sum_{u \in Z \times_Y v} -\dim \mathrm{tot}\phi_u(\mathcal{F}, g \circ f) \\ &= \sum_{u \in Z \times_Y v} (CC\mathcal{F}, d(g \circ f))_{T^*X,u} = (f_* CC\mathcal{F}, dg)_{T^*Y,v} \end{aligned}$$

and the assertion follows. □

If $Y$ is a curve, Conjecture 1.2 may be rephrased as follows. Let $C = SS\mathcal{F}$ be the singular support and assume that on a dense open subscheme $V \subset Y$, the restriction $f_V \colon X_V = X \times_Y V \to V$ of $f$ is $C$-transversal. Then, $f_\circ C \times_Y V$ is a subset of the 0-section. Thus the condition that every irreducible component of $f_\circ C$ is of dimension 1 is satisfied. Further $f_V$ is locally acyclic relatively to $\mathcal{F}$. Since $f$ is proper, $Rf_*\mathcal{F}$ is locally constant on $V$ by [1, Théorème 2.1] and we have

$$CCRf_*\mathcal{F} = -\Big(\mathrm{rank}\ Rf_*\mathcal{F} \cdot [T^*_Y Y] + \sum_{y \in Y - V} a_y Rf_*\mathcal{F} \cdot [T^*_y Y]\Big). \tag{12}$$

For a closed point $y \in Y$, the Artin conductor $a_y Rf_*\mathcal{F}$ is defined by

$$a_y Rf_*\mathcal{F} = \chi(X_{\bar{\eta}}, \mathcal{F}) - \chi(X_{\bar{y}}, \mathcal{F}) + \mathrm{Sw}_y H^*(X_{\bar{\eta}}, \mathcal{F}). \tag{13}$$

In the right hand side, the first two terms denote the Euler-Poincaré characteristics of the geometric generic fiber and the geometric closed fiber respectively and the last term denotes the Swan conductor at $y$.

Let $df$ denote the section of $T^*X$ on a neighborhood of the inverse image $X_y$ defined by the pull-back of a basis $dt$ of the line bundle $T^*Y$ for a local coordinate $t$ on a neighborhood of $y \in Y$. Then, the intersection product $(CC\mathcal{F}, df)_{T^*X,y}$ supported on the inverse image of $X_y$ is well-defined since $SS\mathcal{F}$ is a closed conical subset.

**Lemma 3.** *Let $C = SS\mathcal{F}$ be the singular support and let $V \subset Y$ be a dense open subset such that $f_V \colon X_V \to V$ is projective, smooth and $C$-transversal.*

1. *The equality* (10) *is equivalent to the equality*

$$-a_y Rf_*\mathcal{F} = (CC\mathcal{F}, df)_{T^*X,y} \tag{14}$$

*at each point $y \in Y - V$, where the right hand side denotes the intersection number supported on the inverse image of $y$.*

2. *Further, if $f$ has at most isolated characteristic points, then Conjecture* 1.2 *holds. In particular, if $f \colon X \to Y$ is a finite flat generically étale morphism of smooth curves, then Conjecture* 1.2 *holds.*

3. *Let $\delta_y$ denote the difference of* (14)*. If $X$ and $Y$ are projective, we have $\sum_{y \in Y - V} \delta_y \cdot \deg y = 0$.*

*Proof.* 1. Let $V' \subset V$ be the complement of the images of irreducible components $C_a$ of the singular support $C = \bigcup_a C_a$ such that the image of $C_a$ is a closed point of $Y$. Then, for every closed point $w \in V'$, the immersion $i_w \colon X_w \to X$ is properly $C$-transversal and we have $CCi_w^*\mathcal{F} = i_w^! CC\mathcal{F}$ by [4, Theorem 7.6]. Further, we have

$$f_*CC\mathcal{F} = -\Big((i_w^! CC\mathcal{F}, T^*_{X_w}X_w) \cdot [T^*_Y Y] - \bigcup_{y \in Y - V} (CC\mathcal{F}, df)_{T^*X,y} \cdot [T^*_y Y]\Big). \tag{15}$$

If we assume that $f_V \colon X \times_Y V \to V$ is projective, the index formula (11) implies

$$\operatorname{rank} Rf_*\mathcal{F} = \chi(X_w, i_w^*\mathcal{F}) = (CCi_w^*\mathcal{F}, T^*_{X_w}X_w)_{T^*X_w}.$$

Thus, it suffices to compare (15) and (12).

2. If $f$ has at most isolated characteristic points, (14) is an immediate consequence of the Milnor formula (1).

3. We have

$$\begin{aligned}(CCRf_*\mathcal{F}, T^*_Y Y)_{T^*Y} &= \chi(Y, Rf_*\mathcal{F}) \\ &= \chi(X, \mathcal{F}) = (CC\mathcal{F}, T^*_X X)_{T^*X} = (f_*CC\mathcal{F}, T^*_Y Y)_{T^*Y}\end{aligned}$$

by the index formula (11) and the projection formula. Thus it follows from 1. □

We prove some cases of Conjecture 1.2 assuming that $X$ is a surface.

Let $X$ be a normal noetherian scheme and $U \subset X$ be a dense open subscheme. Let $G$ be a finite group and $V \to U$ be a $G$-torsor. The normalization $Y \to X$ in $V$ carries a natural action of $G$. For a geometric point $\bar{x}$ of $X$, the stabilizer $I \subset G$ of a geometric point $\bar{y}$ of $Y$ above $\bar{x}$ is called an inertia subgroup at $\bar{x}$.

**Lemma 4.** *Let $G$ be a finite group and*

$$\begin{array}{ccc} W & \xleftarrow{j'} & V \\ {\scriptstyle r}\downarrow & & \downarrow{\scriptstyle r'} \\ X & \xleftarrow{j} & U \end{array} \tag{16}$$

*be a cartesian diagram of smooth schemes over a field $k$ where the horizontal arrows are dense open immersions, the right vertical arrow $V \to U$ is a $G$-torsor and the left vertical arrow $r \colon W \to X$ is proper. Assume that for every geometric point $x$ of $X$, the order of the inertia group $I_x \subset G$ is prime to $\ell$.*

*Let $\mathcal{F}$ be a locally constant sheaf on $U$ such that the pull-back $r'^*\mathcal{F}$ is a constant sheaf. Then, for the intermediate extension $j_{!*}\mathcal{F} = j_{!*}(\mathcal{F}[\dim U])[-\dim U]$ on $X$, we have an inclusion*

$$SSj_{!*}\mathcal{F} \subset r_\circ(T^*_W W).$$

*Proof.* Since the assertion is étale local on $X$, we may assume that $G = I_x$ is of order prime to $\ell$. Then, the canonical morphism $\mathcal{F} = (r'_* r'^*\mathcal{F})^G \to r'_* r'^*\mathcal{F}$ is a splitting injection and induces a splitting injection $j_{!*}\mathcal{F} \to j_{!*}r'_* r'^*\mathcal{F}$. Hence, we have

$$SS(j_{!*}\mathcal{F}) \subset SS(j_{!*}r'_* r'^*\mathcal{F}).$$

Since every irreducible subquotient of the shifted perverse sheaf $j_{!*}r'_*r'^*\mathcal{F}$ is isomorphic to an irreducible subquotient of a shifted perverse sheaf ${}^p\mathcal{H}^0(Rr_*j'_*r'^*\mathcal{F}[\dim U])[-\dim U]$ extending $r'_*r'^*\mathcal{F}$, we have

$$SS(j_{!*}r'_*r'^*\mathcal{F}) \subset SS({}^p\mathcal{H}^0(Rr_*j'_*r'^*\mathcal{F})) \subset SS(Rr'_*j'_*r'^*\mathcal{F})$$

by [2, Theorem 1.4 (ii)]. Since $j'_*r'^*\mathcal{F}$ is a constant sheaf on $W$ and $r$ is proper, we have

$$SS(Rr_*j'_*r'^*\mathcal{F}) \subset r_\circ SS(j'_*r'^*\mathcal{F}) \subset r_\circ(T^*_W W)$$

by [2, Lemma 2.2 (ii), Lemma 2.1 (iii)]. Thus the assertion follows. □

**Proposition 5.** *Let $X$ be a normal scheme of finite type over a perfect field $k$, $Y$ be a smooth curve over $k$ and $f\colon X \to Y$ be a flat morphism over $k$. Let $V \subset Y$ be a dense open subscheme such that $f_V\colon X_V = X \times_Y V \to V$ is smooth.*

1. *There exist a finite flat surjective morphism $g\colon Y' \to Y$ of smooth curves over $k$ and a dense open subscheme $X'' \subset X'$ of the normalization $X'$ of $X \times_Y Y'$ satisfying the following condition:*

(1) *We have inclusions $X''_V = X' \times_Y V \subset X'' \subset X'$ and $X'' \subset X'$ is dense in every fiber of $X' \to Y'$. The morphism $X'' \to Y'$ is smooth.*

2. *Let $\mathcal{F}$ be a perverse sheaf on $X_V$ and let $C \subset T^*X_V$ be a closed conical subset on which $\mathcal{F}$ is micro-supported. Assume that $f_V\colon X_V \to V$ is $C$-transversal. Then, there exist $g\colon Y' \to Y$ and $X'' \subset X' \to X \times_Y Y'$ as in* 1. *satisfying the condition* (1) *above and the following condition:*

(2) *Let $j'\colon X''_V \to X''$ denote the open immersion and $C' = SSj'_{!*}\mathcal{F}'$ be the singular support of the intermediate extension $j'_{!*}\mathcal{F}'$ of the pull-back $\mathcal{F}'$ of $\mathcal{F}$ to $X''_V$. Then, the morphism $X'' \to Y'$ is $C'$-transversal.*

*Proof.* By devissage and approximation, we may assume that the complement $Y - V$ consists of a single closed point $y$ and that the closed fiber $X_y$ is irreducible. The assertion is local on a neighborhood in $X$ of the generic point $\xi$ of $X_y$.

1. It follows from [3].

2. Since $f_V\colon X_V \to V$ is smooth, the $C$-transversality of $f_V$ and the condition that $\mathcal{F}$ is micro-supported on $C$ are preserved after base change by [4, Lemma 3.9.2, Lemma 4.2.4]. After replacing $Y$ by $Y'$ and $X$ by $X''$ as in 1., we may assume that $X \to Y$ is smooth. Shrinking $X$ and $Y$ further if necessary, we may assume that $\mathcal{F}$ is locally constant.

Let $W_V \to X_V$ be a $G$-torsor for a finite group $G$ such that the pull-back of $\mathcal{F}$ on $W_V$ is a constant sheaf. Let $r\colon W \to X$ be the normalization of $X$ in $W_V$. Applying 1 to $W \to Y$ and shrinking $X$ if necessary, we may assume that there exists a finite flat surjective morphism of smooth curves $Y' \to Y$ such that the normalization $W'$ of $W \times_Y Y'$ is smooth over $Y'$.

Let $r'\colon W' \to X'$ be the canonical morphism. Since the ramification index at the generic point $\xi'$ of an irreducible component of the fiber $X'_y$ is 1, the inertia group at $\xi'$ is of order a power of $p$. Hence, after shrinking $X$ if necessary, we may assume that for every geometric point $w'$ of $W'$, the order of the inertia group is a power of $p$. Hence, by Lemma 4, we have $C' = SS(j'_{!*}\mathcal{F}) \subset r'_\circ(T^*_{W'}W')$. Since $W' \to Y'$ is smooth, the morphism $f'\colon X' \to Y'$ is $C'$-transversal by [4, Lemma 3.9.3 (2)⇒(1)]. □

**Theorem 6.** *Let the notation be as in Conjecture* 1 *and let $C = SS\mathcal{F}$ be the singular support. Assume that* $\dim X = 2$, $\dim Y = 1$ *and that there exists a dense open subscheme $V \subset Y$ such that $f_V\colon X_V \to V$ is smooth and $C$-transversal. Then, Conjecture* 1.2 *holds.*

*Proof.* We may assume $\mathcal{F}$ is a perverse sheaf by [2, Theorem 1.4 (ii)]. Since the resolution of singularity is known for curves and surfaces, we may assume $Y$ is projective. Since a proper smooth surface over a field is projective, the surface $X$ is projective. Let $y \in Y - V$ be a point. It suffices to show the equality (14).

By Proposition 5 and approximation, there exists a finite flat surjective morphism $Y' \to Y$ of proper smooth curves étale at $y$ and satisfying the conditions in Proposition 5 on the complement $Y - \{y\}$. Since the normalization $X'$ of $X \times_Y Y'$ is projective, we may take a projective smooth scheme $P$ and decompose $f' \colon X' \to Y$ as a composition $X' \to P \to Y$ of a closed immersion $i \colon X' \to P$ and $g \colon P \to Y$.

Let $U = V \cup \{y\}$. Let $\mathcal{F}'$ be the pull-back of $\mathcal{F}$ to $X'_U = X' \times_Y U$ and let $j'_{!*}\mathcal{F}'$ be the intermediate extension with respect to the open immersion $j' \colon X'_U \to X'$. It suffices to show Conjecture 1.2 holds for $P \to Y$ and $\mathcal{G} = i_* j'_{!*}\mathcal{F}'$. Outside the inverse image of $y$, the morphism $g \colon P \to Y$ has at most isolated characteristic points with respect to the singular support $SS\mathcal{G}$ by the condition (2) in Proposition 5 and [4, Lemma 3.9.3 (2)⇒(1)] applied to the restriction of the immersion $X' \to P$ on the complement of a finite closed subset of $X'$. Thus, we have $\delta_{y'} = 0$ for any closed point $y' \in Y'$ not on $y$ by Lemma 3.2. This implies $[Y' \colon Y] \cdot \delta_y = 0$ by Lemma 3.3. Thus the assertion follows. □

**Theorem 7.** *Let the notation be as in Conjecture* 1 *and let $C = SS\mathcal{F}$ be the singular support. Assume that* $\dim X = \dim Y = 2$*, that $f \colon X \to Y$ is proper surjective and that every irreducible component of $f_\circ C$ is of dimension* 2*. Then, Conjecture* 1.2 *holds.*

*Proof.* By Lemma 2, the assertion holds except possibly for the coefficients of the fibers $T^*_y Y$ of finitely many closed points $y \in Y$ where $X \to Y$ is not finite. Let $v \in Y$ be a closed point and we show that the coefficients of the fibers $T^*_v Y$ are equal.

Since the resolution of singularity is known for surfaces and since a proper smooth surface over a field is projective, we may assume that $Y$ and hence $X$ are projective. By replacing $X$ by the Stein factorization of $X \to Y$ except on a neighborhood of $v$, we define $X \to X' \to Y$ such that $f' \colon X' \to Y$ is finite on the complement of $v$ and $r \colon X \to X'$ is an isomorphism on the inverse image of a neighborhood of $v$.

Since $X'$ is projective, we may take a projective smooth scheme $P$ and decompose $f' \colon X' \to Y$ as a composition $X' \to P \to Y$ of a closed immersion $i \colon X' \to P$ and $g \colon P \to Y$. Conjecture 1.2 holds for $\mathcal{G} = i_* Rr_* \mathcal{F}$ and $g \colon P \to Y$ except possibly for the coefficients of the fiber $T^*_v Y$ by Lemma 2. Namely, we have $g_* CC\mathcal{G} = CCRg_*\mathcal{G} = CCRf_*\mathcal{F}$ except possibly for the coefficients of the fiber $T^*_v Y$. By the index formula (11), we have

$$\begin{aligned}(CCRf_*\mathcal{F}, T^*_Y Y)_{T^*Y} &= \chi(Y, Rf_*\mathcal{F}) \\ &= \chi(P, \mathcal{G}) = (CC\mathcal{G}, T^*_P P)_{T^*P} = (g_* CC\mathcal{G}, T^*_Y Y)_{T^*Y}.\end{aligned}$$

Thus, we have an equality also for the coefficients of the fiber $T^*_v Y$. □

We give a characterization of characteristic cycle using functoriality. For the definition of $C$-transversal morphisms $h \colon W \to X$ of smooth schemes and the pull-back $Z_n(C) \to Z_m(h^\circ C)$, we refer to [4, Definition 7.1]. For a constructible complex $\mathcal{F}$ on a projective space $\mathbf{P} = \mathbf{P}^n$, let $R\mathcal{F} = Rp^\vee_* p^* \mathcal{F}[n-1]$ denote the Radon transform on the dual projective space $\mathbf{P}^\vee$ where $p \colon Q \to \mathbf{P}$ and $p^\vee \colon Q \to \mathbf{P}^\vee$ denote the projections on the universal family of hyperplanes $Q = \{(x, H) \in \mathbf{P} \times \mathbf{P}^\vee \mid x \in H\}$. For a linear combination $A = \sum_a m_a C_a$ of irreducible closed conical subset $C_a \subset T^*\mathbf{P}$ of dimension $n$, let $LA = (-1)^{n-1} p^\vee_* p^! A$ denote the Legendre transform (cf. [4, Corollary 7.5]).

**Proposition 8.** *Let $k$ be a perfect field and $\Lambda$ be a finite field of characteristic $\ell$ invertible in $k$. Then, there exists a unique way to attach a linear combination $A(\mathcal{F}) = \sum_a m_a C_a$ satisfying the conditions* (1)-(5) *below of irreducible components of the singular support $SS\mathcal{F} = C = \bigcup_a C_a \subset T^*X$ to each smooth scheme $X$ over $k$ and each constructible complex $\mathcal{F}$ of $\Lambda$-modules on $X$:*

(1) *For every étale morphism $j\colon U \to X$, we have $A(j^*\mathcal{F}) = j^*A(\mathcal{F})$.*

(2) *For every properly $C$-transversal closed immersion $i\colon W \to X$ of smooth schemes, we have $A(i^*\mathcal{F}) = i^!A(\mathcal{F})$.*

(3) *For every closed immersion $i\colon X \to P$ of smooth schemes, we have $A(i_*\mathcal{F}) = i_*A(\mathcal{F})$.*

(4) *For the Radon transform, we have $A(R\mathcal{F}) = LA(\mathcal{F})$.*

(5) *For $X = \operatorname{Spec} k$, we have $A(\mathcal{F}) = \operatorname{rank} \mathcal{F} \cdot T^*_X X$.*

*If $A(\mathcal{F})$ satisfies the conditions* (1)–(5)*, then we have*

$$A(\mathcal{F}) = CC\mathcal{F}. \tag{17}$$

As the proof below shows, it suffices to assume the condition (2) in the case where $i\colon x \to X$ is the closed immersion of a closed point or $i\colon L \to \mathbf{P}$ is the closed immersions of lines in projective spaces. By the definition of the naive Radon transform and the naive Legendre transform, the equality in condition (4) can be decomposed as $A(Rp^\vee_* p^*\mathcal{F}) = p^\vee_* A(p^*\mathcal{F}) = p^\vee_* p^! A(\mathcal{F})$. The first (resp. second) equality corresponds to a special case of Conjecture 1.2 by [4, Lemma 3.11] (resp. to a special case of [4, Proposition 5.17]).

*Proof.* The characteristic cycles satisfy the conditions (1)–(5) by [4, Lemma 5.11.2], [4, Theorem 6.6], [4, Lemma 5.13.2], [4, Corollary 7.12], [4, Lemma 5.11.1] respectively. Thus the existence is proved.

We show the uniqueness. It suffices to show the equality (17). By the condition (2) applied to the closed immersion $x \to X$ of a closed point in a dense open subscheme $U \subset X$ where $\mathcal{F}$ is locally constant and by the conditions (1) and (5), the coefficient of the 0-section $T^*_X X$ in $A(\mathcal{F})$ equals the rank of the restriction $\mathcal{F}|_U$.

We show that this property and the conditions (3)–(5) imply the index formula for projective smooth scheme $X$. By (3), we may assume that $X$ is a projective space $\mathbf{P} = \mathbf{P}^n$ for $n \geqq 2$. Let $R^\vee\mathcal{G} = Rp_*p^{\vee *}\mathcal{G}(n-1)[n-1]$ and $L^\vee B = (-1)^{n-1}p_*p^{\vee !}B$ denote the inverse Radon transform and the inverse Legendre transform. Then, $R^\vee R\mathcal{F}$ is isomorphic to $\mathcal{F}$ up to locally constant complex of rank $(n-2)\cdot\chi(\mathbf{P}_{\bar{k}}, \mathcal{F})$. Hence the coefficient of the 0-section $T^*_{\mathbf{P}}\mathbf{P}$ in $A(R^\vee R\mathcal{F})$ equals that in $A(\mathcal{F})$ plus $(n-1)\cdot\chi(\mathbf{P}_{\bar{k}}, \mathcal{F})$. Similarly, the coefficient of $T^*_{\mathbf{P}}\mathbf{P}$ in $L^\vee LA(\mathcal{F})$ equals that in $A(\mathcal{F})$ plus $(n-1)\cdot(A(\mathcal{F}), T^*_{\mathbf{P}}\mathbf{P})_{T^*\mathbf{P}}$. Thus, we obtain the index formula $\chi(\mathbf{P}_{\bar{k}}, \mathcal{F}) = (A(\mathcal{F}), T^*_{\mathbf{P}}\mathbf{P})_{T^*\mathbf{P}}$ for projective $X$.

Next, we show (17) assuming $\dim X = 1$. Let $U \subset X$ be a dense open subscheme where $\mathcal{F}$ is locally constant. Then, we have

$$A(\mathcal{F}) = -\Big(\operatorname{rank} \mathcal{F} \cdot T^*_X X + \sum_{x \in X - U} m_x \cdot T^*_x X\Big).$$

We show $m_x$ equals the Artin conductor $a_x\mathcal{F}$. If $\mathcal{F}|_U$ is unramified at $x$ and if $\mathcal{F}_x = 0$, the same argument as above shows $m_x = \operatorname{rank} \mathcal{F}|_U = a_x\mathcal{F}$. We show the general case. By (1), we may assume $U = X - \{x\}$. Further shrinking $X$, we may assume that there exists a finite étale surjective morphism $\pi\colon X' \to X$ such that the pull-back $\pi^*\mathcal{F}$ is unramified

at every point $\bar{X}' - X'$ of the boundary of a smooth compactification $j' \colon X' \to \bar{X}'$. Then, we have

$$A(j'_*\pi^*\mathcal{F}) = -\Big(\text{rank } \mathcal{F}|_U \cdot T^*_{\bar{X}'}\bar{X}' + \sum_{x' \in \pi^{-1}(x)} m_x \cdot T^*_{x'}\bar{X}' + \sum_{x' \in \bar{X}' - X'} a_x\mathcal{F} \cdot T^*_{x'}\bar{X}'\Big).$$

Thus, the index formula implies $m_x = a_x\mathcal{F}$.

We show the general case. By (1), we may assume $X$ is affine. We consider an immersion $X \to \mathbf{A}^n \subset \mathbf{P}^n$. Then, by (1) and (3), we may assume $X$ is projective. Set $SS\mathcal{F} = C = \bigcup_a C_a$ and take a projective embedding $X \to \mathbf{P}$ and a pencil $L \subset \mathbf{P}^\vee$ satisfying the following properties as in [5, Lemma 2.3]: The axis $A_L$ of the pencil meets $X$ transversely, that the blow-up $\pi_L \colon X_L \to X$ is $C$-transversal, that the morphism $p_L \colon X_L \to L$ defined by the pencil has at most isolated characteristic points, that the isolated characteristic points are not contained in the inverse image of $V$ and are unique in the fibers of $p_L$, and that for each irreducible component $C_a$ there exists an isolated characteristic point $u$ where a section $dp_L$ of $T^*X$ meets $C_a$.

Then, for $v = p_L(u)$, the coefficient $m_v$ of $T^*_v L$ in $A(Rp_{L*}\pi^*\mathcal{F})$ equals the Artin conductor $-\dim \text{tot}\phi_u(\mathcal{F}, p_L) = -a_v Rp_{L*}\pi^*\mathcal{F}$. Let $i_L \colon L \to \mathbf{P}^\vee$ denote the immersion. Then, by the proper base change theorem and by the conditions (2) and (3), we have $A(Rp_{L*}\pi^*\mathcal{F}) = (-1)^{n-1} i_L^! A(R\mathcal{F}) = i_L^! L A(\mathcal{F})$. Thus, if $A(\mathcal{F}) = \sum_a m_a C_a$, the coefficient $m_v$ equals $m_a \cdot (C_a, dp_L)_{T^*X,u} \neq 0$ and we obtain

$$-\dim \text{tot}\phi_u(\mathcal{F}, p_L) = m_a \cdot (C_a, dp_L)_{T^*X,u} \neq 0.$$

This means that the coefficient $m_a$ is characterized by the same condition as the Milnor formula (1) and we have (17). □

# References


[1] M. Artin, *Théorème de changement de base par un morphisme lisse, et applications*, SGA 4 Exposé XVI, Théorie des Topos et Cohomologie Étale des Schémas, Lecture Notes in Math. 305, 1973, pp 206-249.

[2] A. Beilinson, *Constructible sheaves are holonomic*, `arXiv:1505.06768` to appear at Selecta Math.

[3] H. P. Epp, *Eliminating wild ramification*, Invent. Math. 19 (1973), 235-249.

[4] T. Saito, *The characteristic cycle and the singular support of a constructible sheaf*, `arXiv:1510.03018`, to appear at Inventiones Math.

[5] T. Saito, Y. Yatagawa, *Wild ramification determines the characteristic cycle*, `arXiv:1604.01513`, preprint.